

# Validation of German Aortic Valve Score in a Multi-Surgeon Single Center

Mehmet Kalender<sup>1</sup>, MD; Ahmet Nihat Baysal<sup>1</sup>, MD; Okay Guven Karaca<sup>2</sup>, MD; Kamil Boyacioglu<sup>3</sup>, MD; Nihan Kayalar<sup>3</sup>, MD



DOI: 10.21470/1678-9741-2016-0029

## Abstract

**Objective:** Risk assessment for operative mortality is mandatory for all cardiac operations. For some operation types such as aortic valve repair, EuroSCORE II overestimates the mortality rate and a new scoring system (German AV score) has been developed for a more accurate assessment of operative risk. In this study, we aimed to validate German Aortic Valve Score in our clinic in patients undergoing isolated aortic valve replacement.

**Methods:** A total of 35 patients who underwent isolated open aortic valve replacement between 2010 and 2013 were included. Patients with concomitant procedures and transcatheter aortic valve implantation were excluded. Patients' data were collected and analyzed retrospectively. Patients' risk scores EuroSCORE II were calculated online according to criteria described by EuroSCORE taskforce, Aortic Valve Scores were also calculated.

**Results:** The mean age of patients was 61.14±13.25 years (range 29-80 years). The number of female patients was 14 (40%) and body mass index of 25 (71.43%) patients was in range of 22-35. Mean German Aortic Valve Score was 1.05±0.96 (min: 0 max: 4.98) and mean EuroSCORE was 2.30±2.60 (min: 0.62, max: 2.30). The Aortic Valve Score scale showed better discriminative capacity (AUC 0.647, 95% CI 0.439-0.854). The goodness of fit was  $\chi^2_{HL}[\text{Aortic Valve Score}] = 16.63$ ;  $P = 0.436$ . EuroSCORE II scale had shown less discriminative capacity (AUC 0.397, 95% CI 0.200-0.597). The goodness of fit was good for both scales. The goodness of fit was  $\chi^2_{HL}[\text{EuroSCORE II}] = 30.10$ ;  $P = 0.610$ .

**Conclusion:** In conclusion, German AV score applies to our population with high predictive accuracy and goodness of fit.

**Keywords:** Aortic Valve. Risk Assessment. Adult. Risk Grade.

## Abbreviations, acronyms & symbols

BMI	= Body mass index
EuroSCORE	= European System for Cardiac Operation Risk Evaluation
ROC	= Receiver operating characteristic
TAVI	= Transcatheter aortic valve implantation

## INTRODUCTION

The assessment of operative mortality risk is mandatory for all cardiac operations. Patients need to be informed preoperatively about the risk factors. Some risk scoring systems are used to compare and standardize the results of the operations. The European System for Cardiac Operation Risk Evaluation

(EuroSCORE) is a risk model published in 1999<sup>[1]</sup>. For more than a decade, this risk model had been used widely and validated in innumerable papers demonstrating wonderful goodness of fit<sup>[2,3]</sup>. Current requirements necessitated an update to scoring systems which ended up developing EuroSCORE II which was published on May 2010<sup>[2]</sup>. EuroSCORE II also demonstrated a discriminative capacity similar to EuroSCORE (AUC<sub>EuroSCORE II</sub> = 0.81 vs. AUC<sub>EuroSCORE</sub> = 0.78), and good calibration ( $\chi^2_{HL}[\text{EuroSCORE II}] = 15.48$ ;  $P = 0.0505$ )<sup>[4]</sup>. On the other hand, for specific operation types such as aortic valve repair, EuroSCORE II overestimates the mortality rate<sup>[5-7]</sup> which resulted in development of a new scoring system. Some of these new scoring systems emerged nation based such as Ambler, Guaragna and German Aortic Valve score (formerly named AKL-score)<sup>[8-10]</sup>. German Aortic Valve Score was described by Kötting et al.<sup>[10]</sup> in 2013 with a study in which 1147 isolated aortic valve surgery and transcatheter aortic valve implantation

<sup>1</sup>Konya Education Research Hospital, Cardiovascular Surgery Department, Konya, Turkey.

<sup>2</sup>Duzce University Medical School Hospital, Cardiovascular Surgery Department, Duzce, Turkey.

<sup>3</sup>Bagcilar Education and Research Hospital, Cardiovascular Surgery Department, Istanbul, Turkey.

This study was carried out at the Konya Education Research Hospital, Cardiovascular Surgery Department, Konya, Turkey.

No financial support.

No conflict of interest.

Correspondence Address:

Mehmet Kalender

Cardiovascular Surgery Department, Konya Education and Research Hospital  
Haci saban Mah. Meram Yeniyol cad. No. 92. Konya Eğitim ve Araştırma Hastanesi.  
Meram-Konya, Turkey.

E-mail: ka97084@yahoo.com

Article received on October 2<sup>nd</sup>, 2016.  
Article accepted on December 29<sup>th</sup>, 2016.

(TAVI) patients were enrolled. German aortic valve score has 15 risk factors (Table 1). Two of them (body mass index – BMI – and no sinus rhythm) are different from EuroSCORE II. EuroSCORE II differs in five parameters comparing to German Aortic Valve score (hand poor mobility, diabetes on insulin, Canadian Cardiovascular Society class 4 angina, weight of the intervention and thoracic aorta surgery) – Table 2.

In this study, we aimed to validate German Aortic Valve Score by comparing it with original the EuroSCORE II risk scoring system in patients with isolated open aortic valve replacement.

## METHODS

Patients who underwent isolated open aortic valve replacement between May 2010 and June 2013 were included in the study. Those with concomitant procedures, isolated bioprosthesis replaced patients and TAVI were excluded. Patients'

data were collected and analyzed retrospectively. Primary end point was observed in hospital mortality. Patients' risk scores EuroSCORE II were calculated online according to criteria described by EuroSCORE taskforce<sup>[11]</sup>. Aortic Valve Scores were calculated according to criteria described by Kötting et al.<sup>[10]</sup>.

Sensitivity and specificity was assessed by the use of receiver operating characteristic (ROC) curve and the calibration of German Aortic Valve Score was assessed by Hosmer-Lemeshow (HL) test<sup>[12]</sup>. Calibration was considered to be poor if the test was significant. The discrimination measures the capacity of a model (in this case German Aortic Valve Score and EuroSCORE II) to differentiate the individuals of a sample that suffer an event (in this case, death) and those that do not. The discriminative capacity of the analyzed event was estimated by mean of ROC curve<sup>[13]</sup>. For the analysis, the statistical package SPSS® 15.0 (SPSS, Inc., Chicago, IL, USA) for Windows® was used. A *P*-value <0.05 was considered significant.

**Table 1.** Patients' characteristics. German Aortic Valve Score.

		n	%	Mortality
Age group (years)	<66	20	57.14	5
	66-70	5	14.29	0
	71-75	7	20	1
	76-80	3	8.571	0
Sex	Male	21	60	5
	Female	14	40	1
BMI	22-35	25	71.43	4
	<22	8	22.86	2
	>35	2	5.71	0
Heart failure: NYHA IV	NYHA<IV	34	97.14	6
	NYHA=IV	1	2.85	0
Myocardial infarction < 3 weeks		0	0	0
Critical preoperative status		0	0	0
Pulmonary hypertension		13	37.14	3
No sinus rhythm		4	11.43	1
LVEF (%)	<30	1	2.857	0
	30-50	10	28.57	1
	>50	24	68.57	5
Endocarditis		1	2.85	0
Reoperation		1	2.85	0
Peripheral arterial disease		0	0	0
Chronic obstructive pulmonary disease		5	14.29	1
Chronic renal insufficiency		2	5.714	1
Emergency		2	5.714	0
Observed mortality		6	17.14	6

BMI = body mass index; LVEF = left ventricular ejection fraction; NYHA = New York Heart Association

**Table 2.** Patients' characteristics. EuroSCORE II.

	<b>n</b>	<b>%</b>	<b>Mortality</b>
<b>Patient related factors</b>			
Age (years)		61.14±13.25	6
Female	14	40	1
Peripheral arteriopathy	0	0	0
Chronic obstructive pulmonary disease	5	14.3	1
Diabetes on insulin	3	8.6	0
Poor mobility	0	0	0
<b>Renal impairment</b>			
Dialysis	2	5.71	1
CC<50	5	14.28	0
85<CC>50	20	57.14	2
CC>85	8	22.85	2
<b>Cardiac related factors</b>			
Active endocarditis	1	2.9	0
Recent AMI	0	0	0
<b>NYHA class</b>			
II	31	88.4	4
III	3	8.7	2
IV	1	2.9	0
<b>CCS4</b>	0	0	
<b>LVEF (%)</b>			
>50	24	68.57	5
31-50	10	28.57	1
21-30	1	2.85	0
<20	0	0	0
<b>Pulmonary artery pressure</b>			
31-55 mmHg	3	8.7	0
>55 mmHg	1	2.9	1
<b>Procedure</b>			
Critical Condition	0	0	0
Re-operation	1	2.9	1
Thoracic aorta	0	0	0
<b>Emergency</b>			
Urgent	0	0	0
Emergent	2	5.7	0
Salvage	0	0	0
<b>Weight of procedure</b>			
Single non-CABG	35	100	6

AMI = acute myocardial infarction; CABG = coronary artery bypass grafting; CCS = Canadian Cardiovascular Society; LVEF = left ventricular ejection fraction; NYHA = New York Heart Association

**RESULTS**

We evaluated 35 isolated aortic valve replacement operations in adult patients for this study. The mean age of patients was 61.14±13.25 years (range 29-80 years). The number of female patients was 14 (40%). Patients' characteristics are shown in Tables 1 and 2.

Mean German Aortic Valve Score was 1.05±0.96 (min: 0, max: 4.98) and mean EuroSCORE was 2.30±2.60 (min: 0.62, max: 2.30). The Aortic Valve Score scale showed better discriminative capacity (AUC 0.647, 95% CI 0.439-0.854) (Figure 1). The goodness of fit was  $\chi^2_{HL}[Aortic\ Valve\ Score]=16.63; P=0.436$  (Table 3). EuroSCORE II scale had shown less discriminative capacity (AUC


0.397, 95% CI 0.200-0.597) (Figure 2). The goodness of fit was good for both scales. The goodness of fit was  $\chi^2_{HL}[EuroSCORE\ II]=30.10; P=0.610$  (Table 4).

**DISCUSSION**

Risk scoring systems are valuable for benchmarking of institution results, however, several risk scoring systems have been developed and used. EuroSCORE II is a new updated scoring system with better mortality score and goodness of fit. But some statistical questions have been raised recently<sup>[14,15]</sup>. Moreover, parallel to our opinion there are papers advocating that one scoring system for all patient groups, cardiac diseases


**Table 3.** Contingency table for Hosmer–Lemeshow test (German Aortic Valve Score).

	Observed mortality = 0		Observed mortality = 1		Total
	Observed	Expected	Observed	Expected	Observed
1	4	3.967	0	33	4
2	4	3.778	0	222	4
3	4	4.423	1	577	5
4	4	4.916	2	1.084	6
5	5	3.939	0	1.061	5
6	4	4.514	2	1.486	6
7	4	3.462	1	1.538	5



Diagonal segments are produced by ties.

**Fig. 1 -** The receiver operating characteristic (ROC) curve of German aortic valve score.



Diagonal segments are produced by ties.

**Fig. 2 -** The receiver operating characteristic (ROC) curve of EuroSCORE II.

**Table 4.** Contingency table for Hosmer–Lemeshow test (EuroSCORE II).

	Observed mortality = 0		Observed mortality = 1		Total Observed
	Observed	Expected	Observed	Expected	
1	4	3.972	0	0.028	4
2	4	3.763	0	0.237	4
3	4	3.633	0	0.367	4
4	4	3.443	0	0.557	4
5	1	3.258	3	0.742	4
6	3	3.145	1	0.855	4
7	3	3.003	1	0.997	4
8	4	3.458	1	1.542	5
9	2	1.326	0	0.674	2

and therapies can certainly be misleading<sup>[10,16-18]</sup>. EuroSCORE II was also based on a data set consisting mainly of coronary procedures. Therefore, we believe that there is a requirement for a new scoring system more adaptive for aortic valve procedures. There are also papers reporting the requirement of a new scoring system for aortic valve procedures<sup>[8,10,19-21]</sup>. Kotting et al.<sup>[10]</sup> described a new scoring system for aortic valve procedures based on German Registry.

Former predictive models were developed for specific locations [Ambler, Quaragna Kotting, EuroSCORE and STS], but global need made EuroSCORE and STS popular and they were used widely. As Casalino et al.<sup>[22]</sup> reported in their study that German Aortic Valve Score best fits in German population, but in our opinion it may be applicable to our population as well. Our results showed a high quality of discrimination AUC 0.647 and

Hosmer-Lemeshow method exhibited sufficient concordance in the predicted and observed mortality ( $\chi^2_{HL}$ [Aortic Valve Score]=16.63; P=0.436).

Non-randomized and retrospectively design, single institution setting, multi-surgeon operations and small sample size were the major limitations of our study.

## CONCLUSION

In conclusion, German Aortic Valve score applies to our population with high predictive accuracy and goodness of fit.

## ACKNOWLEDGEMENTS

Special thanks to Necla Yildiz and Cemile Cevik for their enormous help during the writing process.

## Authors' roles & responsibilities

MK	Conception and study design; realization of operations; analysis and/or data interpretation; statistical analysis; manuscript redaction or critical review of its content; final manuscript approval
ANB	Conception and study design; realization of operations; analysis and/or data interpretation; statistical analysis; manuscript redaction or critical review of its content; final manuscript approval
OGK	Conception and study design; realization of operations; analysis and/or data interpretation; statistical analysis; manuscript redaction or critical review of its content; final manuscript approval
KB	Conception and study design; realization of operations; analysis and/or data interpretation; statistical analysis; manuscript redaction or critical review of its content; final manuscript approval
NK	Conception and study design; realization of operations; analysis and/or data interpretation; statistical analysis; manuscript redaction or critical review of its content; final manuscript approval

## REFERENCES

1. Roques F, Nashef SA, Michel P, Gauducheau E, de Vincentiis C, Baudet E, et al. Risk factors and outcome in European cardiac surgery: analysis of the EuroSCORE multinational database of 19030 patients. *Eur J Cardiothorac Surg.* 1999;15(6):816-23.
2. Roques F, Michel P, Goldstone A, Nashef SA. The logistic EuroSCORE. *Eur Heart J.* 2003;24(9):881-2.
3. Siregar S, Groenwold RH, de Heer F, Bots ML, van der Graaf Y, van Herwerden LA. Performance of the original EuroSCORE. *Eur J Cardiothorac Surg.* 2012;41(4):746-54.
4. Nashef SA, Roques F, Sharples LD, Nilsson J, Smith C, Goldstone AR, et al. EuroSCORE II. *Eur J Cardiothorac Surg.* 2012;41(4): 734-44.
5. Gummert JF, Funkat A, Osswald B, Beckmann A, Schiller W, Krian A, et al. EuroSCORE overestimates the risk of cardiac surgery: results from the national registry of the German Society of Thoracic and Cardiovascular Surgery. *Clin Res Cardiol.* 2009;98(6):363-9.
6. Biancari F, Kangasniemi OP, Aliasim Mahar M, Rasinaho E, Satomaa A, Tiozzo V, et al. Changing risk of patients undergoing coronary artery bypass surgery. *Interact Cardiovasc Thorac Surg.* 2009;8(1):40-4.
7. Nilsson J, Algotsson L, Höglund P, Lühns C, Brandt J. Comparison of 19 pre-operative risk stratification models in open-heart surgery. *Eur Heart J.* 2006;27(7):867-74.

8. Ambler G, Omar RZ, Royston P, Kinsman R, Keogh BE, Taylor KM. Generic, simple risk stratification model for heart valve surgery. *Circulation*. 2005;112(2):224-31.
9. Guaragna JC, Bodanese LC, Bueno FL, Goldani MA. Proposed preoperative risk score for patients candidate to cardiac valve surgery. *Arq Bras Cardiol*. 2010;94(4):541-8.
10. Kötting J, Schillerb W, Beckmann A, Schäfer E, Döbler K, Hamm C, et al. German Aortic Valve Score: a new scoring system for prediction of mortality related to aortic valve procedures in adults. *Eur J Cardiothorac Surg*. 2013;43(5):971-7.
11. EuroSCORE. Available at <http://www.euroscore.org/> Accessed March 21, 2014.
12. Hosmer DW, Lemeshow S. A goodness-of-fit test for the multiple logistic regression model. *Comm in Stat*. 1980;A10:1043-69.
13. Royston P, Moons KG, Altman DG, Vergouwe Y. Prognosis and prognostic research: developing a prognostic model. *BMJ*. 2009;338:b604.
14. Sergeant P, Meuris B, Pettinari M. EuroSCORE II, illum qui est gravitates magni observe. *Eur J Cardiothorac Surg*. 2012;41(4):729-31.
15. Collins GS, Altman DG. Design flaws in EuroSCORE II. *Eur J Cardiothorac Surg*. 2013;43(4):871.
16. Hannan EL, Wu C, Bennett EV, Carlson RE, Culliford AT, Gold JP, et al. Risk index for predicting in-hospital mortality for cardiac valve surgery. *Ann Thorac Surg*. 2007;83(3):921-9.
17. Zheng Z, Li Y, Zhang S, Hu S, Chinese CABG Registry Study. The Chinese coronary artery bypass grafting registry study: how well does the EuroSCORE predict operative risk for Chinese population? *Eur J Cardiothorac Surg*. 2009;35(1):54-8.
18. Howell NJ, Head SJ, Freemantle N, van der Meulen TA, Senanayake E, Menon A, et al. The new EuroSCORE II does not improve prediction of mortality in high-risk patients undergoing cardiac surgery: a collaborative analysis of two European centres. *Eur J Cardiothorac Surg*. 2013;44(6):1006-11.
19. Florath I, Rosendahl UP, Mortasawi A, Bauer SF, Dalladaku F, Ennker IC, et al. Current determinants of operative mortality in 1400 patients requiring aortic valve replacement. *Ann Thorac Surg*. 2003;76(1):75-83.
20. Jin R, Grunkemeier GL, Starr A, Providence Health System Cardiovascular Study Group. Validation and refinement of mortality risk models for heart valve surgery. *Ann Thorac Surg*. 2005;80(2):471-9.
21. Nowicki ER, Birkmeyer NJ, Weintraub RW, Leavitt BJ, Sanders JH, Dacey LJ, et al.; Northern New England Cardiovascular Disease Study Group and the Center for Evaluative Clinical Sciences, Dartmouth Medical School. Multivariable prediction of in-hospital mortality associated with aortic and mitral valve surgery in Northern New England. *Ann Thorac Surg*. 2004;77(6):1966-77.
22. Casalino R, Tarasoutchi F, Spina G, Katz M, Bacelar A, Sampaio R, et al. EuroSCORE models in a cohort of patients with valvular heart disease and a high prevalence of rheumatic fever submitted to surgical procedures. *PLoS One*. 2015;10(2):e0118357.